



Forest Service  
U.S. DEPARTMENT OF AGRICULTURE

National Office | GTR WO-99 | December 2021

# Forests to Faucets 2.0

## Connecting Forests, Water, and Communities





# Forests to Faucets 2.0

## Connecting Forests, Water, and Communities

**Erika Mack**  
Resource Information Specialist  
*U.S. Department of Agriculture, Forest Service,  
Southern Research Station, Eastern Forest  
Environmental Threat Assessment Center*  
3041 East Cornwallis Road  
Research Triangle Park, NC 27709

**Ge Sun**  
Research Hydrologist and Project Leader  
*U.S. Department of Agriculture, Forest Service,  
Southern Research Station, Eastern Forest  
Environmental Threat Assessment Center*  
3041 East Cornwallis Road  
Research Triangle Park, NC 27709

**Rebecca Lilja**  
Geographic Information System Specialist  
*U.S. Department of Agriculture, Forest Service,  
State and Private Forestry, Eastern Region*  
271 Mast Road  
Durham, NH 03824

**Peter Caldwell**  
Research Hydrologist  
*U.S. Department of Agriculture, Forest Service,  
Southern Research Station, Center for Integrated  
Forest Science*  
3160 Coweeta Lab Road  
Otto, NC 28763

**Sally Claggett**  
Liaison to the Chesapeake Bay Program  
*U.S. Department of Agriculture, Forest Service,  
State and Private Forestry, Eastern Region*  
410 Severn Avenue, Suite 209  
Annapolis, MD 21403

### ABSTRACT

The Forest to Faucets version 2.0 (F2FV2) assessment uses geospatial modeling at the 12-digit Hydrologic Unit Code (HUC12) scale in the conterminous United States to identify watersheds that are most important to surface drinking water, the ability to produce clean water, forest ownership (public or private), and potential threats to water yield from insects and diseases, wildfire, land use or climate change. F2FV2 updates a 2011 version of the project (Forests to Faucets version 1.0). Results, presented by regions administered by the U.S. Department of Agriculture, Forest Service, indicate that watersheds in the Eastern, Southern, and Pacific Southwest Regions were most important for surface drinking water. Watersheds in the Southern, Pacific Northwest, and Pacific Southwest Regions had the highest ability to produce clean water based on the five biophysical characteristics evaluated. The Pacific Southwest, Pacific Northwest, and Northern Regions had the most watersheds at the highest threat of wildfire as well as the most watersheds at the highest threat of insects and disease. For all future climate and population growth scenarios, the Southern, Pacific Southwest, and Eastern Regions had the most watersheds at the highest threat of land use change, while the Pacific Northwest and Southern Regions had the most watersheds at the highest threat of decreases in water yield because of climate change. F2FV2 provides a user-friendly tool and relatively high spatial resolution benchmark dataset that forest managers can use to evaluate the effect of their management on the water supply, and that water consumers can use for considering potential threats upstream.

**Keywords:** climate change, land use change, drinking water, forest, insects and disease, surface water, water, water demand, watershed, wildfire

### Contents

**ABSTRACT . . . . .**

**INTRODUCTION . . . . .**

Threats to Surface Drinking Water . . . . .

“Forests to Faucets” Assessments . . . . .

**METHODS. . . . .**

Extent and Scale of Analysis. . . . .

Models. . . . .

Ability To Produce Clean Water (APCW) Index Model . . . . .

Threats to Surface Drinking Water  
(THREAT) Model . . . . .

Data . . . . .

Population and Surface Drinking . . . . .

Water Intakes. . . . .

Water Yield . . . . .

Land Cover . . . . .

Riparian Area. . . . .

Land Use Change . . . . .

Insects and Disease. . . . .

Wildfire . . . . .

Forest Ownership . . . . .

**RESULTS . . . . .**

Index of Importance to Surface Drinking Water (IMP) . . . . .

Ability to Produce Clean Water (APCW) Index . . . . .

Threats to Surface Drinking Water . . . . .

Water Yield Decrease . . . . .

Land Use Change . . . . .

Insect and Disease . . . . .

Wildfire . . . . .

**SUMMARY. . . . .**

**LIST OF ABBREVIATIONS. . . . .**

**REFERENCES. . . . .**

**2**

**4**

**5**

**6**

**7**

**7**

**9**

**10**

**11**

**12**

**12**

**12**

**13**

**14**

**14**

**15**

**16**

**16**

**16**

**17**

**17**

**17**

**19**

**19**

**22**

**22**

**26**

**28**

**29**

**30**





USDA Forest Service photo by Joshua Courter.

## INTRODUCTION

Forests are an important source of clean drinking water for over 150 million people in the United States (Caldwell et al. 2014; Liu et al. 2020). The 749 million acres of forested lands provide more than half of the national water yield from lands in the lower 48 States (Sun et al. 2015; Liu et al., 2021). Approximately 74 percent of total drinking water withdrawals originates from surface water sources, such as streams, ponds, and reservoirs (Dieter et al. 2018). However, there are increasing concerns about the quantity and quality of both ground water and surface water supplies because of climate change, population growth, changes in land use, and increased water demand (Sun et al. 2008).

### Threats to Surface Drinking Water

Disturbances that interrupt core aspects of the hydrologic cycle pose threats to surface drinking water. In particular, land use change, insects and disease, wind events, and wildfire can reduce or eliminate vegetative cover (i.e., trees and shrubs) and ground cover (i.e., litter, impervious surfaces). The loss of vegetation reduces rainfall interception, water infiltration, and evapotranspiration, which can increase the amount of overland flow, stormflow, and streamflow, respectively, and can also alter the timing of that flow. Land use change and increases in flow can lead to a decrease in water quality from non-point source pollutants through increased nutrient and sediment loads and often toxins, heavy metals, and other chemicals (e.g., from towns and cities, farming practices, fire retardant, and burned infrastructure). When the core aspects of

the hydrologic cycle are altered, the overall status of the watershed condition degrades (Hallema et al. 2018, 2019; Neary et al. 2005; Sun and Caldwell 2015).

Similarly, changes in climate can alter the magnitude, timing, and quality of surface water supplies. Temperature and precipitation are the two key climatic variables that control water yield (i.e., runoff). Changes in these variables—specifically higher temperatures and increased variability of precipitation—threaten forests directly by changing their structure and functions, and key ecosystem services such as clean water supply. Indirect threats to forests related to climate change include more frequent and intense wildfires and insect and disease outbreaks under prolonged drought stress and high temperatures (Koch and Coulston 2020; Vose et al. 2012, 2018; Westerling et al. 2006) that affect watershed health and water quantity and quality.



“Forests to Faucets” Assessments

Land and water resource managers need comprehensive data and tools for planning and management of watersheds threatened by land use change, climate change, wildfire, and insects and disease. In response to these needs, the U.S. Department of Agriculture (USDA), Forest Service, State and Private Forestry completed the “From the Forest to the Faucet: Drinking Water and Forests in the U.S.” version 1.0 (F2FV1) assessment (Weidner and Todd 2011). The F2FV1 assessment had multiple objectives that served land managers on a broad scale, including (Weidner and Todd 2011):

1. Using geospatial analysis at the 12-digit Hydrologic Unit Code (HUC12) scale to identify watersheds in the conterminous United States that are most important to surface drinking water, the forests protecting surface drinking water in those watersheds, and forested areas that face threats from increased housing density insects and disease, and the frequency and severity of wildfires.
2. Creating a dataset that can be useful for identifying priority areas for protecting the quality of surface drinking water and can incorporate into broad-scale planning efforts or existing decision support tools.
3. Identifying watersheds that could be targeted in a payment-for-watershed-services project.
4. Creating an educational tool that highlights the role of forests in providing an ecosystem service of surface drinking water.

For the F2FV1 assessment, Weidner and Todd (2011) used four geospatial models:

1. Index of importance to surface drinking water (IMP) model.
2. Drinking water protection (PR) model.
3. Index of forest importance to surface drinking water (FIMP) model.
4. Index of threats to forest importance to surface drinking water model.

The assessment found that many watersheds in the Eastern United States had higher importance to surface

drinking water than those in the West and Midwest. The assessment also showed how forests—often privately owned—in watersheds of high importance were key to the production of surface drinking water, but the threat assessment of water quality showed that the ability of forests in these watersheds to sustain important surface drinking water supplies was at threat. The results of the assessment were informative and produced multiple map layers that stakeholders could easily integrate into decision support tools (Weidner and Todd 2011). For example, the State of Vermont incorporated the F2FV1 assessment into its 2017 forest action plan to show areas of the highest priority for protecting and conserving public drinking water supplies (Vermont Department of Forests, Parks, and Recreation 2017).

Forests to Faucets version 2.0 (F2FV2) expands on the F2FV1 assessment, with updates to the methodology and input data. The F2FV2 project had multiple objectives and new features, including:

1. Using geospatial analysis at the HUC12 scale for the conterminous United States to determine watersheds most important to surface drinking water; type of forest ownership; and where they are threatened by insects and disease, wildfire, land use change, or climate change that decreases water yield.
2. Identifying watersheds with the natural ability to produce clean water under current land use conditions.
3. Developing an interactive tool for stakeholders to access and use the results of this work.

This report focuses on the methods the authors of this document used for completing the assessment and provides an overview of the results, which can be downloaded or viewed from the Forests to Faucets website: [https://www.fs.fed.us/ecosystems-services/FS\\_Efforts/forests2faucets.shtml](https://www.fs.fed.us/ecosystems-services/FS_Efforts/forests2faucets.shtml).

METHODS

Extent and Scale of Analysis

The F2FV2 study focused on watershed characteristics and water supply and demand for the conterminous United States at the HUC12 resolution (10,000 to 40,000 acres, or about 40 square kilometers [km<sup>2</sup>]

to 160 km<sup>2</sup>).<sup>1</sup> The 2019 version of the Watershed Boundary Dataset (WBD) (U.S. Department of Agriculture, Natural Resources Conservation Service 2019) provided the HUC12 vector delineation for the analysis and the watershed connectivity. This project considered 83,314 HUC12s within the conterminous United States, and 8 administrative geographic regions of the Forest Service summarized the results in this report (figure 1):

- Northern Region
- Rocky Mountain Region
- Southwestern Region
- Intermountain Region
- Pacific Southwest Region
- Pacific Northwest Region
- Southern Region
- Eastern Region

Most of the data used for the conterminous United States analysis of F2FV2 were not available for Alaska, Hawaii, and the territories of Puerto Rico, the U.S. Virgin Islands, Guam, the Mariana Islands, and American Samoa. However, land cover and the U.S. Environmental Protection Agency (EPA) Safe Drinking Water Information System (SDWIS) intake data were available, and used to calculate the risk-based drinking water protection (PR) model for these areas. These outputs are available at the Forests to Faucets website [www.fs.fed.us/ecosystems-services/FS\\_Efforts/forests2faucets.shtml](http://www.fs.fed.us/ecosystems-services/FS_Efforts/forests2faucets.shtml).



Figure 1—The administrative regions of the Forest Service within the conterminous United States.

1. A hydrologic unit is “a topographically defined set of drainage areas organized in a nested hierarchy by size and number of divisions per nested level” (U.S. Department of the Interior, Geological Survey and U.S. Department of Agriculture, Natural Resources Conservation Service 2013).



Models

The authors used four models in the development of F2FV2: the IMP model, the PR model, the ability to produce clean water (APCW) index model, and the threats to surface drinking water (THREAT) model (table 1). The F2FV2 assessment used the same model equations from the F2FV1 assessment, with updated input data, to calculate the IMP and PR indices. Weidner and Todd (2011) did not use the APCW model in the F2FV1 assessment; Barnes et al. (2009) describe the foundation of this model. This assessment uses a modified version of the APCW model that included five attributes to characterize the biophysical conditions within a watershed. The F2FV2 THREAT model differs from the index of threats to forest importance to surface drinking water model that Weidner and Todd (2011) used in the F2FV1 assessment; F2FV used a different equation to calculate threats and added climate change to the list of threats assessed.

Index of Importance to Surface Drinking Water (IMP) Model and Drinking Water Protection (PR) Model

The authors compiled water yield (i.e., water supply) and the population served by surface water intakes (water demand) to estimate the IMP index for watershed n. IMP was calculated using equation (1):

Model	Equation	Parameters
Index of Importance to Surface Drinking Water (IMP) model	$IMP_n = Q_n \times PR_n$	$Q_n$ is average annual water yield (mm/year) $PR_n$ is the Drinking Water Protection model
Drinking Water Protection (PR) model	$PR_n = \sum(W_i \times P_i)$	$W_i$ is a proportional weighting $P_i$ is the total population served by intakes in the $i$ th watershed downstream from watershed $n$
Ability to Produce Clean Water (APCW) Index model	$APCW = (N + A + I + R) \times Q$	$N$ is percentage of natural cover $A$ is percentage of agricultural land $I$ is percentage of impervious surface $R$ is percentage of riparian natural cover $Q$ is average annual water yield (mm/year)
Threats to Surface Drinking Water (THREAT) model	$THREAT = (IMP \times APCW \times PT) / 10,000$	IMP is the Index of Importance to Surface Drinking Water model APCW is the Ability to Produce Clean Water Index model PT is percent of a watershed that is threatened from a potential threat to surface water. PT is defined differently for each threat, climate change that decreases water yield ( $PT_{YIELD}$ ), land use change ( $PT_{LUC}$ ), insects and disease ( $PT_{ID}$ ), or wildfire potential ( $PT_{WFP}$ )

Table 1—Models used in the Forests to Faucets version 2.0 assessment

IMP\_n = Q\_n x PR\_n

where

Q\_n = the average annual water yield in millimeters (mm) per year for each watershed n

PR\_n = the drinking water protection model (water demand calculation for watershed n)

PR\_n was determined using equation (2):

PR\_n = Σ(W\_i x P\_i)

where

W\_i = the proportional weighting.

P\_i = the total population served by intakes in the ith watershed downstream from watershed n (Weidner and Todd 2011) (figure 2).

The PR model identifies the importance of a watershed’s land area by using the downstream drinking water demand. Watersheds with the highest water demand are considered the most important. The population served by surface water intakes located within the watershed represents water demand in the model. Through watershed connectivity, the model accounts for water demand within the source watershed, plus a fraction of water demand for all watersheds downstream




Figure 2—Schematic diagram illustrating the calculation of IMP Index where Q is average annual runoff (mm/year), P<sub>i</sub> is population being served by the drinking water intake (green dot), and W<sub>i</sub> is the proportional weight, which represents the fraction of water demand from the downstream watersheds that contribute to the importance of the source watershed and is expressed by an exponential decay relationship.



of the source watershed. The proportional weighting,  $W_i$ , expressed by an exponential decay relationship, represents the fraction of water demand from the downstream watersheds that contribute to the importance of the source watershed.  $W_i$  was calculated using equation (3):

$$W_i = (1 - 0.01)^d$$

where  
 $d$  = the distance, assumed to be 25 kilometers (km) in stream length, between the source watershed and a downstream watershed containing an intake (table 2) (Weidner and Todd 2011). Inter-basin transfers or water diversions were not incorporated into the model.

In the F2FV1 assessment, Weidner and Todd (2011) derived the exponential decay relationship equation after reviewing literature about decay curves from a variety of contaminants and consulting with a science advisory team. The equation is meant to represent, in a generalized way, the decay of contaminants in a stream or river (Weidner and Todd 2011). The final IMP value was divided into 10 quantiles and mapped on a 0 to 100 scale.

**Ability To Produce Clean Water (APCW) Index Model**  
This assessment used land cover and water yield datasets to calculate the APCW index, a water quality index that reflects watershed integrity by incorporating the conditions of five attributes. APCW characterizes the



USDA Forest Service photo.

biophysical conditions that relate to the ability to provide clean water in each watershed. APCW was calculated using equation (4):

$$APCW = (N + A + I + R) \times Q$$

where  
 $N$  = percentage of natural cover (ranking points).  
 $A$  = percentage of agricultural land (ranking points).  
 $I$  = percentage of impervious surface (ranking points).  
 $R$  = percentage of riparian area that is natural cover (ranking points).  
 $Q$  = the average annual water yield in mm per year (ranking points).

i <sup>th</sup> downstream HUC	Distance down-stream ( <i>d</i> ) (km)	Proportional weight ( <i>W<sub>i</sub></i> )
0	0	1.000
1	25	0.779
2	50	0.607
3	75	0.472
4	100	0.368
5	125	0.287
6	150	0.223
7	175	0.174
8	200	0.135
9	225	0.105
10	250	0.082

Table 2—Proportional weight values used in the Index of Importance to Surface Drinking Water (IMP) model

This assessment ranks the attributes from low to very high, based on the percentage of that attribute in a watershed, and values the attributes at 1 to 4 points, respectively (table 3), based on accepted standards in the scientific literature (Barnes et al. 2009). Point values of the four attributes were summed and multiplied by the ranked average annual water yield (table 3). The APCW index was determined by dividing the results by the maximum score of 64. APCW values were divided into 10 quantiles and mapped on a 0 to 100 scale.

**Threats to Surface Drinking Water (THREAT) Mode**  
Combining the IMP and APCW indices with each surface drinking water potential threat (PT) produced the THREAT index for each watershed.. THREAT, an environmental threat index, was calculated using equation (5):

$$THREAT = (IMP \times APCW \times PT) / 10,000$$

where  
 $IMP$  = index of importance to surface drinking water model.  
 $APCW$  = ability to produce clean water index model.  
 $PT$  = percent of a watershed that is threatened from a potential threat to surface water.

PT is defined differently for each threat: climate change that decreases water yield ( $PT_{YIELD}$ ), land use change ( $PT_{LUC}$ ), insects and disease ( $PT_{ID}$ ), or wildfire potential ( $PT_{WFP}$ ). Storm events such as hurricanes, tornadoes, and straight-line wind events were considered but were not used because no national coverage dataset was available. THREAT values were divided by 10,000 to maintain a 0 to 100 index.

Attribute	Very high (4 points)	High (3 points)	Moderate (2 points)	Low (1 point)
N = percentage of natural cover	>75	51–75	25–50	<25
A = percentage of agricultural land	<10	10–20	21–30	>30
I = percentage of impervious surface	0–1	2–5	6–10	>10
R = percentage of riparian natural cover	>70	51–70	30–50	<30
Q = average annual water yield (mm/year)	>800	401–800	201–400	0–200

Table 3—The Ability to Produce Clean Water Index model attribute ranking system

$PT_{YIELD}$  was calculated for low emissions and high emissions future climate scenarios and for future years 2040 and 2090 using equations (6–9):

$$PT_{YIELD(LOW40)} = [((Yield2010 - Yield2040low) / (Yield2010)) \times 100]$$

$$PT_{YIELD(LOW90)} = [((Yield2010 - Yield2090low) / (Yield2010)) \times 100]$$

$$PT_{YIELD(HIGH40)} = [((Yield2010 - Yield2040high) / (Yield2010)) \times 100]$$

$$PT_{YIELD(HIGH90)} = [((Yield2010 - Yield2040high) / (Yield2010)) \times 100]$$

where  
Yield2010, Yield2040, and Yield2090 are the water yield as calculated by the Water Supply Stress Index (WaSSI) model for the years 2010, 2040, and 2090 respectively by a 20-year average around the year of interest.

This study only considered watersheds that saw a decrease in water yield as potential threat from climate change. While the duration and timing of precipitation events can pose threats to watersheds with no water yield change and infrastructure can be potentially threatened by increases in water yield, detailed examinations of these factors were not considered in this analysis. Refer to the data section for a detailed description of the climate datasets and the WaSSI model.



$PT_{LUC}$  was calculated for a low emissions and high emission future climate scenarios and for future years 2040 and 2090 using equations (10–13):

$$PT_{LUC(LOW40)} = (LUC2010 - LUC2040_{low}) / Total\_HUC12$$

$$PT_{LUC(LOW90)} = (LUC2010 - LUC2040_{low}) / Total\_HUC12$$

$$PT_{LUC(HIGH40)} = (LUC2010 - LUC2090_{high}) / Total\_HUC12$$

$$PT_{LUC(HIGH90)} = (LUC2010 - LUC2090_{high}) / Total\_HUC12$$

Where LUC2010, LUC2040, and LUC2090 are the land use change as calculated by the Integrated Climate and Land-Use Scenarios (ICLUS) for the years 2010, 2040, and 2090, and Total\_HUC12 is the total HUC12 area for the HUC12 of interest. The change that occurred was one direction, i.e., from a nonurban land use classification to an urban classification. Refer to the data section for a detailed description of the land use datasets.

$PT_{ID}$  was calculated using equation (14):

$$PT_{ID} = (At\_Risk / TOTAL\_ID) \times 100$$

where At\_Risk is the treed area having National Insect and Disease Risk Map (NIDRM)-modelled “risk” and TOTAL\_ID is the sum of the treed area having NIDRM-modelled “risk” and “not at risk.” Refer to the data section for a description of the insect and disease dataset.

$PT$  was calculated using equation (15):

$$PT_{WFP} = (Total\_H\_VH / TOTAL\_WFP) \times 100$$

where Total\_H\_VH is the sum of the watershed area in the wildfire potential classes of high and very high; TOTAL\_WFP is the sum of watershed area in all the other classes present in wildfire potential dataset. Refer to the data section for a description of the wildfire potential dataset.

2. The Great Lakes present a special case in that there are a large number of drinking water intakes found offshore. To maintain a similar weighting scenario using the population served by each water intake, the authors assigned offshore intakes to the watershed within 25 kilometers of the intake and used them together with all the on-land intakes in the drinking water protection (PR) model. Because intakes are located offshore, all areas bordering the Great Lakes affect the water quality of the offshore intakes, not just the nearest subwatersheds where they were assigned.

Data

Inputs for the IMP, APCW, and THREAT models (table 4) used a variety of national scale data. The authors rescaled the data used in the assessment models from their native resolution to the HUC12 boundaries.

**Population and Surface Drinking Water Intakes**  
This study used the EPA’s SDWIS dataset (U.S. Environmental Protection Agency 2018) to identify drinking water intake locations and the population served by those intakes for the IMP model. SDWIS contains basic, violation, and enforcement information on water systems in the United States. The States self-report the data contained in SDWIS to the EPA, and the EPA aggregates these to a national dataset as submitted (without correction). The authors used basic information from SDWIS, including the water system identification (ID), the latitude and longitude of the water system intake, the population served by the water system, and the source water type (surface water, ground water, etc.)This analysis omitted intakes with erroneous location information and only considered intakes with a source water type code of surface water or ground water under the direct influence of surface water. The population served by each water intake was calculated by summing the population within a water system and dividing it by the total number of intakes within that water system.<sup>2</sup> The analysis included a total of 16,811 water intakes within the conterminous United States; the total population served by those intakes was 124,835,279. The IMP model used these data as inputs.

**Water Yield**  
The authors used the water supply stress index (WaSSI) model to create the water yield (runoff) dataset by quantifying the amount of water produced within a HUC12 watershed under various climate scenarios. The USDA Forest Service developed WaSSI to assess the potential impacts of changes in population, climate, and land use on water availability and ecosystem productivity at the conterminous United States scale (Caldwell et al. 2012; Lockaby et al. 2013; Marion et al. 2013; Sun et al. 2008, 2011a, 2011b, 2013; Tavernia et al. 2013). WaSSI runs at the monthly time step and comprises water balance, ecosystem productivity, and water supply and demand modules.

The authors averaged baseline water yield as annual

Data	Parameter	Source	Resolution	Time period	Model(s)
Watershed boundary	Watershed area	Watershed Boundary Dataset (WBD) (USGS and USDA NRCS 2013)  ( <a href="https://www.usgs.gov/core-science-systems/ngp/national-hydrography/watershed-boundary-dataset">https://www.usgs.gov/core-science-systems/ngp/national-hydrography/watershed-boundary-dataset</a> )	HUC12	2019	IMP, APCW, THREAT
Population and surface drinking water intakes	Water intake, population served	Safe Drinking Water Information System (SDWIS) database (EPA 2017) ( <a href="https://www.epa.gov/enviro/sdwis-search">https://www.epa.gov/enviro/sdwis-search</a> )	Aggregated to the HUC12 watershed	2017	IMP
Historical climate	Monthly total precipitation, mean temperature	Parameter-elevation Regressions on Independent Slopes Model (PRISM) data (Daly and others 1994) ( <a href="https://prism.oregonstate.edu/">https://prism.oregonstate.edu/</a> )	4 km x 4 km	1960–2015	Used to estimate water yield for IMP, APCW, THREAT
Future climate	Monthly total precipitation, mean temperature	Multivariate Adaptive Constructed Analogs (MACA) datasets (Taylor and others 2012)  ( <a href="https://climate.northwestknowledge.net/MACA/index.php">https://climate.northwestknowledge.net/MACA/index.php</a> )	6 km x 6 km	1961–2099	Used to estimate water yield for IMP, APCW, THREAT
Water Yield	Average annual water yield, percent change in average annual water yield	Water Supply Stress Index (WaSSI) model (Caldwell and others 2012; Lockaby and others 2013; Marion and others 2013; Sun and others 2008, 2011a, 2011b, 2013; Tavernia and others 2013)  ( <a href="https://forestthreats.org/research/tools/WaSSI/">https://forestthreats.org/research/tools/WaSSI/</a> )	HUC12	1960–2099	IMP, APCW, THREAT
Land cover	Developed; forest area; percentage of agriculture, impervious, and natural cover	National Land Cover Database (NLCD) (USGS 2019)  ( <a href="https://www.mrlc.gov/data">https://www.mrlc.gov/data</a> )	30 m x 30 m	2016	APCW
Riparian area	Natural cover	National riparian areas base map (Abood and others 2012a, 2012b) ( <a href="https://www.riparian.solutions/story-map">https://www.riparian.solutions/story-map</a> )	10 m x 10 m	2019	APCW
Land use change	Future land use	Integrated Climate and Land Use Scenarios (ICLUS) version 2.1 (EPA 2017) ( <a href="https://iclus.epa.gov/">https://iclus.epa.gov/</a> )	90 m x 90 m	2000–2100	THREAT
Insect and disease	Insect/disease risk area	National Insect and Disease Risk Map (Krist and others 2014) ( <a href="https://www.fs.fed.us/foresthealth/applied-sciences/mapping-reporting/national-risk-maps.shtml">https://www.fs.fed.us/foresthealth/applied-sciences/mapping-reporting/national-risk-maps.shtml</a> )	240 m x 240 m	2012	THREAT
Wildfire	Wildfire risk area	Forest Service Wildfire Hazard Potential (WHP) dataset (Dillon 2018)  ( <a href="https://www.firelab.org/project/wildfire-hazard-potential">https://www.firelab.org/project/wildfire-hazard-potential</a> )	270 m x 270 m	2018	THREAT
Forest ownership	Forest ownership	National Conservation Easement Database (NCED) version 3 ( <a href="https://databasin.org/datasets/">https://databasin.org/datasets/</a> )	Parcel	2013	None. Ancillary layer available in online tool for filtering

Table 4—Input datasets for the Forests to Faucets version 2.0 assessment. APCW = Ability to Produce Clean Water Index model; IMP = Index of Importance to Surface Drinking Water model; THREAT = Threats to Surface Drinking Water model



water yield for a historical time period (1961 to 2015) and estimated future water yield for two future time periods (2040, an average of the years 2030 to 2049; and 2090, an average of the years 2080 to 2099). The parameter-elevation regression on independent slopes model (PRISM) was the source of the historical climate data (Daly et al. 1994). The HadGEM2-ES365 general circulation model using representative concentration pathways (RCP) greenhouse gas emissions scenarios 4.5 and 8.5 was the source of the future climate data (Taylor et al. 2012). The RCP 4.5 emissions scenario is considered a less warming scenario, and RCP 8.5 is considered a higher warming scenario (Hayhoe et al. 2018). This assessment calculated the percent change in water yield by subtracting the average annual water yield for the historical time period from the future time period and dividing by the historical water yield for each scenario. Inputs for the IMP and APCW models used the average annual water yield; THREAT model inputs used the percent change in average annual water yield.

Land Cover

Every 5 years, the Multi-Resolution Land Characteristics (MRLC) Consortium produces the National Land Cover Database (NLCD), which was used to estimate land cover for the APCW model. The NLCD includes 30- x 30-meter (m) resolution land cover, percentage of developed imperviousness, and land cover change products for the United States. The MRLC uses the Landsat 5 thematic mapper (TM) as the source for the land cover classification; Homer et al. (2015); the U.S. Department of the Interior,

Geological Survey (2019); and Yang et al. (2018) categorized the United States into 16 classes. The 2016 version of the NLCD provided inputs for the following APCW modeling parameters under baseline conditions: percentage of natural cover, percentage of agricultural land, percentage of impervious surface, and percentage of riparian natural cover. The dataset also provided forest cover input for forest ownership. This study grouped together classes of NLCD land cover to delineate these input parameters (table 5).

Riparian Area

The Forest Service developed a 10- x 10-m resolution riparian areas base map dataset for the continental United States and delineated it to the 50-year flood height. This dataset is unique in that it integrates streams, lakes, watersheds, wetlands, soils, elevation, land cover, and hydrologic data to delineate a variable-width riparian area (Abood et al. 2012a, 2012b). The authors overlaid the riparian areas base map on the 2016 NLCD dataset with two classifications—natural cover and developed and agriculture (table 5)—to determine the percentage of the riparian area with natural cover at 30- x 30-m resolution, which was input into the APCW model.

Land Use Change

The THREAT model used the EPA’s Integrated Climate and Land-Use Scenarios (ICLUS) version 2.1 dataset to estimate the threat of land use change across the United States (U.S. Environmental Protection Agency 2017). The ICLUS dataset projects decadal population into the year 2100, based on the Intergovernmental Panel on Climate Change (IPCC)

shared socioeconomic pathways (SSP), which affects fertility, mortality, and immigration. ICLUS uses a mathematical model to simulate population migration patterns at the county level to estimate housing demand. Finally, a statistical model predicts future impervious surface area from the housing demand estimate.

ICLUS creates land use scenarios by combining standard SSPs—possible future socioeconomic conditions with either no climate change for the year 2010 or a climate change projection identified by the IPCC standard RCPs. Climate change projections used two future time periods: 2040 (an average of the years 2030 to 2049) and 2090 (an average of the years 2080 to 2099). This study used six ICLUS datasets (table 6). The combination of SSP5 and RCP8.5 represents a higher emissions, higher population-growth scenario, and the combination of SSP2 and RCP4.5 represents a lower emissions, lower population-growth scenario (U.S. Environmental Protection Agency 2017).

The ICLUS dataset has a 90- x 90-m resolution and 19 land-use classes (table 7). The authors reclassified land-use classes 14 to 18 to the land-use class “urban.” The assessment calculated the land area that changed to “developed” based on the SSP-RCP scenario for each future time period and also calculated the percentage of land changed to “developed” for HUC12 watersheds. The authors input the percentage of a watershed predicted to change to “developed” into the THREAT model for each of the climate emissions scenarios.

Insects and Disease

This study used the 2012 version of the National Insect and Disease Risk Map (NIDRM) to estimate risk of insects and disease in the THREAT model. Every 5 years, the Forest Service, Forest Health Technology Enterprise Team produces the NIDRM for the United States. The NIDRM defines risk as “the expectation that, without remediation, at least 25 percent of standing live basal area greater than 1 inch in diameter will die over a 15-year timeframe (2013 to 2027) due to insects and diseases” (Krist et al. 2014). The dataset has a 240- x 240-m resolution, and the NIDRM modeled 1.2 billion acres of treed areas (“any areas where presence of trees was recorded” [Krist et al. 2014]), as opposed to forested areas obtained from a remotely sensed land cover map that identifies forest location. The authors tabulated the area of the two classes—risk and no risk—within each HUC12 at the 30- x 30-m resolution, and from that, calculated the percentage of HUC12 with NIDRM-modeled risk, which was incorporated into the THREAT model.

Wildfire

The 2018 wildfire hazard potential (WHP) dataset, developed by the Forest Service to “help inform evaluations of wildfire risk” (Dillon 2018), provided input for the wildfire risk parameter. The raster dataset has a 270- x 270-m resolution and covers the conterminous United States. “Areas mapped with higher WHP values represent fuels and other landscape conditions with a higher probability of experiencing high-intensity fire with torching, crowning, and other forms of extreme wildfire behavior under conducive weather conditions”

Input Parameter	Forest to Faucets Definition (NLCD class grouping)	Model
Forest ownership	41, 42, 43, 90	Used as filter in the F2FV2 online tool
Percentage of agricultural land	81, 82	APCW
Percentage of impervious surface	NLCD percent developed imperviousness dataset	APCW
Percentage of natural cover	11, 12, 41, 42, 43, 51, 52, 71, 90, 95	APCW
Percentage of riparian cover	21, 22, 23, 24, 31, 81, 82	APCW

Table 5—Crosswalk of National Land Cover Database (NLCD) land use class grouping to input parameters for the Forests to Faucets version 2.0 (F2FV2) assessment

Shared Socioeconomic Pathways (SSP)	Climate Change Model	Year
SSP2	No climate change	2010
	HadGEM2-ES365 RCP 4.5	2040
	HadGEM2-ES365 RCP 4.5	2090
SSP5	No climate change	2010
	HadGEM2-ES365 RCP 8.5	2040
	HadGEM2-ES365 RCP 8.5	2090

Table 6—The land use and climate scenarios from the Integrated Climate and Land-Use Scenarios (ICLUS) version 2.1 dataset used in the Forests to Faucets version 2.0 assessment



Code	Group	Class name	Forest to Faucets classification
0	Water	Natural water	Not reclassified
1		Reservoirs, canals	Not reclassified
2		Wetlands	Not reclassified
3	Protected	Recreation, conservation	Not reclassified
4	Working/production	Timber	Not reclassified
5		Grazing	Not reclassified
6		Pasture	Not reclassified
7		Cropland	Not reclassified
8		Mining, barren land	Not reclassified
9	Developed	Parks, golf courses	Not reclassified
10		Exurban, low density	Not reclassified
11		Exurban, high density	Not reclassified
12		Suburban	Not reclassified
13		Urban, low density	Not reclassified
14		Urban, high density	Urban
15		Commercial	Urban
16		Industrial	Urban
17		Institutional	Urban
18		Transportation	Urban

Table 7—The Integrated Climate and Land-Use Scenarios (ICLUS) land use classification scheme

(Dillon et al. 2015). The dataset presents WHP in seven classes: very low, low, medium, high, very high, nonburnable, and water. The authors tabulated the area of each WHP class within HUC12 watersheds at the processing resolution for the project—30 m x 30 m—and calculated the percentage of the high and very high WHP classes for each HUC12 watershed. The resulting percentage map provided input for the THREAT model.

Forest Ownership

This study integrated the Conservation Biology Institute’s Protected Areas Database of the United States (PAD-US) version 2.1, the National Conservation Easement Database (NCED) version 3, and NLCD forest land cover classes to determine forest ownership by HUC12 watershed. The timeframe of the forest ownership for the three datasets represents (approximately) 2016. The PAD-US identifies lands that are in fee simple ownership. The NCED identifies lands that have permanent conservation

easements—private lands voluntarily set aside by legal means between the owner and the Government, or a land trust for restricted use to protect the land’s conservation value. The PAD-US and NCED account for approximately 990 million acres of land within the United States (Foster et al. 2014). The authors merged the PAD-US and NCED vector datasets together into a single dataset and grouped them into five categories (table 8). The dataset was then converted to a raster with a 30- x 30-m resolution. Next, the PAD-US and NCED layers were combined with the NLCD forest dataset (classes 41, 42, 43, and 90) to tabulate the area of forest ownership within the HUC12 watersheds. Finally, the assessment calculated the percentage of protected forested land per HUC12 as an ancillary layer for use in the web tool as a filter.

Forests to Faucets class name	PAD-US/NCED class name
Other Federal forest, not national forest	Federal land
National forest	USDA Forest Service
Nonforest	Nonforest
Private forest	Private forest
Protected forest	NCED permanent, Native American land, joint ownership, local land, private conservation land, State land, unknown

Table 8—Crosswalk of Protected Areas Database of the United States (PAD-US)/National Conservation Easement Database (NCED) class to Forests to Faucets version 2.0 ownership class

RESULTS

The following results summarize the F2FV2 model findings for the eight regions within the conterminous United States (figure 1).

Index of Importance to Surface Drinking Water (IMP)

The IMP index is an estimated value that integrates average annual runoff, population, and water intake data for the watersheds. This study maps the index on a scale of 0 to 100 and charted with five classes in 20-unit increments, ranging from very low to very high importance for surface drinking water. The IMP model results identified the Forest Service Eastern, Southern, and Pacific Southwest Regions as those with the most watersheds with very high importance to surface drinking water supplies (figures 3 and 4). The Southwestern and Intermountain regions had the most watersheds with very low importance to surface drinking water supplies (figures 3 and 4). In general, the watersheds with very high importance to surface drinking water supplies are those that correspond to locations with large populations that rely on surface drinking water and have a higher water use. These results are consistent with the results from the F2FV1 assessment (Weidner and Todd 2011).




Figure 3—The Importance to Surface Drinking Water (IMP) Index map for the 83,314 watersheds in the conterminous United States. Watersheds with darker blue colors have higher importance for protecting surface drinking water.






Figure 4—Percentage of a Forest Service region area within an Importance to Surface Drinking Water (IMP) Index class.

## Ability to Produce Clean Water (APCW) Index

The APCW model integrated the percentage of natural cover, percentage of agricultural land, percentage of impervious surface, percentage of riparian natural cover, and average annual runoff of a watershed to determine the APCW index. This study maps the index on a scale of 0 to 100 and charted with five classes in 20-unit increments, ranging from a very low to a very

high ability to produce clean water. The APCW model results identified the Southern, Pacific Northwest, and Pacific Southwest Regions as having the most watersheds with a very high ability to produce clean water (figures 5 and 6). The Rocky Mountain and Northern Regions had the most watersheds with a very low ability to produce clean water (figures 5 and 6). In general, watersheds with a high percentage of vegetation and low percentage of impervious surface would have a higher ability to produce clean water.




Figure 5—The Ability to Produce Clean Water (APCW) Index map for the 83,314 watersheds in the conterminous United States. Watersheds in blue have the highest ability to produce clean water.




Figure 6—Percentage of a Forest Service region area within an Ability to Produce Clean Water (APCW) Index class.



Threats to Surface Drinking Water

The THREAT model provided the framework for determining the threats to surface drinking water from a range of potential threats, including insects and disease, wildfire, climate change that reduces water yield, and land use change. Combining IMP and APCW indices with each quantified potential threat produced a THREAT index for each. The authors mapped these THREAT indices on a scale of 0 to 100 to allow integration with the IMP and APCW map that identified watersheds important to surface drinking water and the ability to produce clean water.

Water Yield Decrease

This assessment considered climate change trends in terms of decreasing water availability that results from changes in water balance due to increasing temperatures and changes in precipitation in U.S.

watersheds. Figure 7 shows watersheds important to drinking water that were predicted by the WaSSI model to experience a decrease in water yield. The Pacific Northwest, Pacific Southwest, and Southern Regions had the most watersheds in the very high THREAT index class under the 2040 low-emissions scenario (figures 7A and 8A), and the Southern, Eastern, and Pacific Northwest Regions had the most watersheds in the very high THREAT index class under the 2040 high-emissions scenario (figures 7C and 8C).

The Southern, Pacific Southwest, and Pacific Northwest Regions had the most watersheds in the very high THREAT index class under the 2090 low-emissions scenario (figures 7B and 8B), and the Southern, Pacific Northwest, and Eastern Regions had the most watersheds in the very high THREAT index class under the 2090 high-emissions scenario (figures 7D and 8D). For all of the future climate scenarios, all




Figure 7—Watersheds important to surface drinking water supply predicted by the WaSSI model to experience a decrease in water yield under future climate scenarios (A and B: low emissions scenarios for 2040 and 2090, respectively; C and D: high emissions scenarios for 2040 and 2090, respectively). Watersheds in red have the highest THREAT index.




Figure 8—Percentage of a Forest Service region area predicted by the WaSSI model to experience a decrease in water yield under future climate scenarios (A and B: low emissions scenarios for 2040 and 2090, respectively; C and D: high emissions scenarios for 2040 and 2090, respectively).

of the regions had some watersheds in the very high THREAT index class, but the Pacific Northwest and Southern Regions consistently had the most watersheds in the very high THREAT index class for the future, and could be priority areas of focus in future work.

Land Use Change

Figure 9 shows watersheds important to surface drinking water that were predicted by ICLUS to have increased land use change due to population growth. The Southern, Pacific Southwest, and Eastern Regions had the most watersheds in the very high THREAT index class for land use change for both the low and high emission scenarios for the 2040 and 2090 time periods (figures 9 and 10). These results reflect historical population trends of the United States, with the Southern United States having the highest rate of population growth since 2000, according to the U.S. Census Bureau (U.S. Department of Commerce, Census Bureau 2019; Wilson et al. 2012).

Insect and Disease

Figure 11 shows watersheds important to surface drinking water that have a threat of tree mortality from insect and disease. The analysis showed 19 percent of all HUC12 watersheds within the conterminous United States had threat from insect and disease. The Northern, Pacific Northwest, and Pacific Southwest Regions had the most watersheds in the very high THREAT index class for insects and disease (figures 11 and 12). In 2016 and 2018, the Forest Service, Forest Health Protection’s National Insect and Disease Survey identified the west coast (the Pacific Northwest and Pacific Southwest Regions) as having the largest area (1.95 million hectares [ha] in 2016 and 1.08 million ha in 2018) with mortality agents and complexes. During 2016 and 2018, the top three mortality agents threatening west coast forests included fir engraver (*Scolytus ventralis*), western pine beetle (*Dendroctonus brevicomis*), and mountain pine beetle (*D. ponderosae*) (Potter et al. 2018, 2020).






Figure 9—Watersheds important to surface drinking water supply predicted by ICLUS to have increased land use change due to population growth under future climate scenarios (A and B: low emissions scenarios for 2040 and 2090, respectively; C and D: high emissions scenarios for 2040 and 2090, respectively). Watersheds in red have the highest THREAT index.




Figure 10—Percentage of a Forest Service region area predicted by ICLUS to have increased land use change under future climate scenarios (A and B: low emissions scenarios for 2040 and 2090, respectively; C and D: high emissions scenarios for 2040 and 2090, respectively).






Figure 11—Watersheds important to surface drinking water supply that have a threat of tree mortality from insects and disease. Watersheds in red have the highest THREAT.




Figure 12—Percentage of a Forest Service region area that have a threat of tree mortality from insects and disease.



Wildfire

Figure 13 shows watersheds important to surface drinking water classified with high or very high wildfire hazard potential, based on the 2018 WHP dataset. Nineteen percent of all HUC12 watersheds had a threat from wildfire, and the remaining eighty-one percent had little to no threat. The Pacific Southwest, Pacific Northwest, and Northern Regions had the most watersheds in the very high THREAT index class for wildfire (figures 13 and 14). In the conterminous

United States, northern California and north-central Washington ecoregions, which overlay the Pacific Southwest and Pacific Northwest Regions, had the highest fire occurrence densities (the number of fire occurrences per 100 km<sup>2</sup> [10,000 ha] of tree canopy coverage area) in 2018. For the years 2001 to 2017, the California, northern Rocky Mountains, Southwest, and Southeastern Coastal Plain ecoregions had the highest annual mean number of fire occurrences per 100 km<sup>2</sup> of tree canopy coverage area (Potter 2020).




Figure 13—Watersheds important to surface drinking water supply at threat from wildfires.




Figure 14—Percentage of a Forest Service region area at threat from wildfire.



SUMMARY

By employing new datasets, Forests to Faucets version 2.0 (FTFV2) represents an update of a previous effort to explicitly map the connections between forests and surface drinking water supply, with a relatively high spatial resolution (HUC12) at a national level. This assessment incorporated new environmental threats such as climate change and land use change scenarios along with detailed land data characteristics (i.e., riparian cover).

When examining the THREAT index results for each Forest Service region, the authors classified most of the watersheds as low or very low. This was expected because of the driving factors for the potential threat considered in this report are localized. For example, an insect and disease outbreak can be specific to a tree species, and that species may not be present in all watersheds within a Forest Service region. The potential for wildfire risk depends on the availability of fuels, and all watersheds within a region may not have a buildup of fuels. All watersheds within a Forest Service region will not see a reduction of water or an increase of population in the future. The THREAT maps are tools land and water resource managers can use to identify potential hotspots that could require further investigation and potential mitigation.

By Forest Service region, the importance of watersheds (IMP index) to surface drinking water and the ability

to produce clean water (APCW index) are more regional and contiguous in nature. The patterns of IMP index are consistent with patterns of stream flow routing that is incorporated within the index methodology; the biophysical characteristics of the APCW index generally span contiguous watersheds.

This study was designed to use commonly available national datasets to model water quantity and quality at the watershed scale. The Forests to Faucets version 2.0 dataset and maps are most appropriate for use at the large basin, and regional to national scale. Thus, land and water resource managers can incorporate the assessment results into decision support tools at the State, regional, or national scale, or in forest management plans. Land and water resource managers should be cautious using the dataset and results solely for making decisions local small scales, as localized data would be more appropriate. Predicting the response of water quantity and quality to future environmental change can be extremely challenging, and the results of this study are subject to validation and revisions. This study provides a tool and benchmark dataset for land and water resource managers to evaluate the impact of their management on surface water supply and for water consumers to consider potential environmental threats upstream, now and in the future.



USDA Forest Service photo.

LIST OF ABBREVIATIONS

APCW	the ability to produce clean water index model
EPA	U.S. Environmental Protection Agency
F2FV1	Forest to Faucets version 1.0
F2FV2	Forest to Faucets version 2.0
FIMP	index of forest importance to surface drinking water model
HUC12	12-digit hydrologic unit code
ICLUS	integrated climate and land-use scenarios
ID	identification
IMP	index of importance to surface drinking water model
IPCC	Intergovernmental Panel on Climate Change
MRLC	Multi-Resolution Land Characteristics
NCED	national conservation easement database
NIDRM	national insect and disease risk map
NLCD	national land cover database
PAD-US	protected areas database of the United States
PR	drinking water protection model
PRISM	parameter-elevation regression on independent slopes model
PT	potential threat
RCP	representative concentration pathways
SDWIS	safe drinking water information system
SSP	shared socioeconomic pathways
THREAT	threats to surface drinking water model
TM	thematic mapper
WaSSI	water supply stress index model
WHP	wildfire hazard potential



## REFERENCES

- Abood, S.; Maclean, A. 2012a. Modeling and classifying riparian ecotones via GIS utilizing geophysical and vegetative inputs: a new approach. In: Lanfear, K.J.; Spangenberg, N.E., eds. AWRA 2012 summer specialty conference: riparian ecosystems IV: advancing science, economics, and policy 2012. Middleburg, VA: American Water Resources Association: 22–27.
- Abood, S.; Maclean, A.; Mason, L. 2012b. Modeling riparian zones utilizing DEMS and flood height data. *Photogrammetric Engineering and Remote Sensing*. 78(3): 259–269. <https://doi.org/10.14358/PERS.78.3.259>.
- Barnes, M.; Todd, A.; Whitney Lilja, R.; Barten, P. 2009. Forests, water and people: drinking water supply and forest lands in the Northeast and Midwest United States, June 2009. NA-FR-01-08. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. 71 p.
- Caldwell, P.V.; Muldoon, C.; Ford-Miniat, C. [et al.]. 2014. Quantifying the role of National Forest System lands in providing surface drinking water supply for the Southern United States. Gen. Tech. Rep. SRS-197. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 135 p. <https://doi.org/10.2737/SRS-GTR-197>.
- Caldwell, P.V.; Sun, G.; McNulty, S.G. [et al.]. 2012. Impacts of impervious cover, water withdrawals, and climate change on river flows in the conterminous US. *Hydrology and Earth System Sciences Discussions*. 9: 4263–4304. <https://doi.org/10.5194/hessd-9-4263-2012>.
- Daly, C.; Neilson, R.P.; Phillips, D.L. 1994. A statistical-topographic model for mapping climatological precipitation over mountainous terrain. *Journal of Applied Meteorology and Climatology*. 33(2): 140–158. [https://doi.org/10.1175/1520-0450\(1994\)033<0140:ASTMFM>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0140:ASTMFM>2.0.CO;2).
- Dieter, C.A.; Maupin, M.A.; Caldwell, R.R. [et al.]. 2018. Estimated use of water in the United States in 2015. U.S. Geological Survey Circular 1441. Reston, VA: U.S. Department of the Interior, Geological Survey. 65 p. <https://doi.org/10.3133/cir1441>.
- Dillon, G.K. 2018. Wildfire Hazard Potential (WHP) for the conterminous United States (270-m GRID), version 2018 classified. 2d ed. Fort Collins, CO: Forest Service Research Data Archive. <https://doi.org/10.2737/RDS-2015-0046-2>.
- Dillon, G.K.; Menakis, J.; Fay, F. 2015. Wildland fire potential: a tool for assessing wildfire risk and fuels management needs. In: Keane, R.E.; Jolly, M.; Parsons, R.; Riley, K., tech. eds. *Proceedings of the large wildland fires conference*. RMRS-P-73. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: 60–76.
- Foster, K.; Comendant, T.; Lundin, M.; Strittholt, J. 2014. Protecting natural vegetation: comparative analysis of land protection mechanisms. Corvallis, OR: Conservation Biology Institute. 31 p.
- Hallema, D.W.; Sun, G.; Caldwell, P.V. [et al.]. 2018. Burned forests impact water supplies. *Nature Communications*. 9: 1307. <https://doi.org/10.1038/s41467-018-03735-6>.
- Hallema, D.W.; Sun, G.; Caldwell, P.V. [et al.]. 2019. Wildland fire impacts on water yield across the contiguous United States. Gen. Tech. Rep. SRS-238. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 109 p. <https://doi.org/10.2737/SRS-GTR-238>.
- Hayhoe, K.; Wuebbles, D.J.; Easterling, D.R. [et al.]. 2018. Our changing climate. In: Reidmiller, D.R.; Avery, C.W.; Easterling, D.R. [et al.], eds. *Fourth national climate assessment: impacts, risks, and adaptation in the United States, volume II*. Washington, DC: U.S. Global Change Research Program: 72–144. Chap. 2. <https://doi.org/10.7930/NCA4.2018.CH2>.
- Homer, C.G.; Dewitz, J.A.; Yang, L. [et al.]. 2015. Completion of the 2011 National Land Cover Database for the conterminous United States: representing a decade of land cover change information. *Photogrammetric Engineering and Remote Sensing*. 81(5): 345–354.
- Koch, F.H.; Coulston, J.W. 2020. Drought and moisture surplus patterns in the conterminous United States: 2018, 2016–2018, and 2014–2018. In: Potter, K.M.; Conkling, B.L., eds. *Forest health monitoring: national status, trends, and analysis 2019*. Gen. Tech. Rep. SRS-250. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 83–102. Chapter 4.
- Krist, F.J., Jr.; Ellenwood, J.R.; Woods, M.E. [et al.]. 2014. 2013–2027 National insect and disease forest risk assessment. FHTET-14-01. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team. 199 p. [https://www.fs.fed.us/foresthealth/technology/pdfs/2012\\_RiskMap\\_Report\\_web.pdf](https://www.fs.fed.us/foresthealth/technology/pdfs/2012_RiskMap_Report_web.pdf).
- Liu, N.; Dobbs, G.R.; Caldwell, P.V. [et al.]. 2020. Quantifying the role of State and private forest lands in providing surface drinking water supply for the Southern United States. Gen. Tech. Rep. SRS-248. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 405 p. <https://doi.org/10.2737/SRS-GTR-248>.
- Lockaby, G.; Nagy, C.; Vose, J. [et al.]. 2013. Forests and water. In: Wear, D.N.; Greis, J.G., eds. *The Southern Forest Futures Project: technical report*. Gen. Tech. Rep. SRS-GTR-178. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station: 309–339.
- Marion, D.A.; Sun, G.; Caldwell, P.V. [et al.]. 2013. Managing forest water quantity and quality under climate change. In: Vose, J.M.; Klepzig, K.D., eds. *Climate Change Adaptation and Mitigation Management Options: a guide for natural resource managers in southern forest ecosystems*. Boca Raton, FL: CRC Press: 249–305.
- Neary, D.G.; Ryan, K.C.; DeBano, L.F. 2005. Wildland fire in ecosystems: effects of fire on soils and water. Gen. Tech. Rep. RMRS-GTR-42. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 250 p. Vol. 4. <https://doi.org/10.2737/RMRS-GTR-42-V4>.
- Potter, K.M. 2020. Broad-scale patterns of forest fire occurrence across the 50 United States, and the Caribbean Territories, 2018. In: Potter, K.M.; Conkling, B.L., eds. *Forest health monitoring: national status, trends, and analysis 2019*. Gen. Tech. Rep. SRS-250. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station: 57-82. Chap. 3.



- Potter, K.M.; Paschke, J.L.; Koch, F.H. 2020. Large-scale patterns of insect and disease activity in the conterminous United States, Alaska, and Hawaii from the national Insect and Disease Survey, 2018. In: Potter, K.M.; Conkling, B.L., eds. Forest health monitoring: national status, trends, and analysis 2019. Gen. Tech. Rep. SRS-250. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station: 27–56. Chap. 2.
- Potter, K.M.; Paschke, J.L.; Zweifler, M.O. 2018. Large-scale patterns of insect and disease activity in the conterminous United States, Alaska, and Hawaii from the national Insect and Disease Survey, 2016. In: Potter, K.M.; Conkling, B.L., eds. Forest health monitoring: national status, trends, and analysis 2017. Gen. Tech. Rep. SRS-233. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 23–44. Chap. 2.
- Sun G.; Alstad, K.; Chen, J. [et al.]. 2011a. A general predictive model for estimating monthly ecosystem evapotranspiration. *Ecohydrology*. 4: 245–255. <https://doi.org/10.1002/eco.194>.
- Sun, G.; Caldwell, P.; Noormets, A. [et al.]. 2011b. Upscaling key ecosystem functions across the conterminous United States by a water-centric ecosystem model. *Journal of Geophysical Research Biogeosciences*. 116: G00J05. <https://doi.org/10.1029/2010JG001573>.
- Sun, G.; Caldwell, P. 2015. Impacts of urbanization on stream water quantity and quality in the United States. *Water Resources Impact*. 17(1): 17–20.
- Sun, G.; Caldwell, P.V.; McNulty, S.G. [et al.]. 2013. Impacts of climate change and variability on water resources in the Southeast. In: Ingram, K.T.; Dow, K.; Carter, L., eds. Southeastern regional technical report to the National Climate Change Assessment. [Place of publication unknown]: Island Press: 204–234. Chap. 10.
- Sun, G.; Caldwell, P.V.; McNulty, S.G. 2015. Modelling the potential role of forest thinning in maintaining water supplies under a changing climate across the conterminous United States. *Hydrological Processes*. 29(24): 5016–5030. <https://doi.org/10.1002/hyp.10469>.
- Sun, G.; McNulty, S.G.; Moore Myers, J.A.; Cohen, E.C. 2008. Impacts of multiple stresses on water demand and supply across the Southeastern United States. *Journal of the American Water Resource Association*. 44(6): 1441–1457. <https://doi.org/10.1111/j.1752-1688.2008.00250.x>.
- Tavernia, B.G.; Nelson, M.D.; Caldwell, P.V.; Sun, G. 2013. Water stress projections in the Northeastern and Midwestern United States in 2060: anthropogenic and ecological consequences. *Journal of the American Water Resources Association*. 49(4): 938–952. <https://doi.org/10.1111/jawr.12075>.
- Taylor, K.E.; Stouffer, R.J.; Meehl, G.A. 2012. An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*. 93(4): 485–498. <https://doi.org/10.1175/BAMS-D-11-00094.1>.
- U.S. Department of Agriculture [USDA], Natural Resources Conservation Service [NRCS]. 2019. Watershed Boundary Dataset for continental United States. <http://www.nrcs.usda.gov/wps/portal/nrcs/main/national/water/watersheds/dataset>. (1 September 2019).
- U.S. Department of Commerce [DOC], Census Bureau. 2019. Annual estimates of the resident population: April 1, 2010 to July 1, 2018; United States metropolitan and micropolitan statistical area, and for Puerto Rico. <https://www2.census.gov/programs-surveys/popest/tables/2010-2018/state/totals/PEPANNRES.pdf>. (23 August 2019).
- U.S. Department of the Interior, Geological Survey [USGS]. 2019. National Land Cover Database (NLCD) 2016 land cover (CONUS). <https://www.mrlc.gov/data>. (14 December 2020).
- U.S. Department of the Interior, Geological Survey [USGS]; U.S. Department of Agriculture [USDA], Natural Resources Conservation Service [NRCS]. 2013. Federal standards and procedures for the national Watershed Boundary Dataset (WBD). 4th ed.: Techniques and Methods 11-A3. Reston, VA: U.S. Geological Survey. 63 p. <https://pubs.usgs.gov/tm/11/a3>. (19 October 2020).
- U.S. Environmental Protection Agency [EPA]. 2017. Updates to the demographic and spatial allocation models to produce Integrated Climate and Land-Use Scenarios (ICLUS) version 2. EPA/600/R-16/366F. Washington, DC: U.S. Environmental Protection Agency. 63 p. plus appendixes.
- U.S. Environmental Protection Agency [EPA]. 2018. Safe Drinking Water Information System. [Online database]. Washington, DC: U.S. Environmental Protection Agency. <https://ofmpub.epa.gov/apex/sfdw/f?p=108:200>. (December 2018).
- Vermont Department of Forests, Parks, and Recreation. 2017. 2017 forest action plan. [https://fpr.vermont.gov/forest/vermonts\\_forests/action\\_plan](https://fpr.vermont.gov/forest/vermonts_forests/action_plan). (5 May 2020).
- Vose, J.M.; Peterson, D.L.; Domke, G.M. [et al.]. 2018. Forests. In: Reidmiller, D.R.; Avery, C.W.; Easterling, D.R. [et al.], eds. Fourth national climate assessment: impacts, risks, and adaptation in the United States, volume II. Washington, DC: U.S. Global Change Research Program: 232–267. Chap. 6. <https://doi.org/10.7930/NCA4.2018.CH6>.
- Vose, J.M.; Peterson, D.L.; Patel-Weynand, T., eds. 2012. Effects of climatic variability and change on forest ecosystems: a comprehensive science synthesis for the U.S. forest sector. Gen. Tech. Rep. PNW-GTR-870. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 265 p. <https://doi.org/10.2737/PNW-GTR-870>.
- Weidner, E.; Todd, A. 2011. From the forest to the faucet: drinking water and forests in the US. Methods paper. Washington, DC: U.S. Department of Agriculture, Forest Service. 34 p.
- Westerling, A.L; Hidalgo, H.G.; Cayan, D.R. [et al.]. 2006. Warming and earlier spring increase Western U.S. forest wildfire activity. *Science*. 313(5789): 940–943. <https://doi.org/10.1126/science.1128834>.
- Wilson, S.G.; Plane, D.A.; Mackun, P.J. [et al.]. 2012. Patterns of metropolitan and micropolitan population change: 2000 to 2010. 2010 Census special reports: C2010SR-01. Washington, DC: U.S. Department of Commerce, Census Bureau. 89 p.
- Yang, L.; Jin, S.; Danielson, P. [et al.]. 2018. A new generation of the United States National Land Cover Database: requirements, research priorities, design, and implementation strategies. *ISPRS Journal of Photogrammetry and Remote Sensing*. 146: 108–123. <https://doi.org/10.1016/j.isprsjprs.2018.09.006>.



In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](#) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

USDA is an equal opportunity provider, employer, and lender.



